

Preface

I present below [the fundamental principles](#) I've derived for making the most of ySTR DNA testing for genealogical purposes.

I qualify “DNA testing” with the technical term [ySTR](#) (short for Y-Chromosome Satellite Repeat) merely to differentiate the two kinds of tests that Family Tree DNA (FTDNA) and others have offered, the other kind being the [ySNP](#) tests which are at present more suited to exploring human population history than genealogy. The definitional links are to the glossary of terms on my DNA testing information page, which provides an introduction to many different types of testing, as well as links to texts and papers relevant to the interpretation of these tests. However, it is my goal here to bring together in one place everything that is germane to the interpretation of a set of ySTR results for a particular [genealogical patrilineage](#).

Besides “patrilineage” there are a number of other terms related to DNA testing for genealogical purposes that it's essential to understand, and for all of these I have provided hyperlinks below at the point where they are first introduced.

The other prefatory point I wish to make is that much of my advice below runs counter to what you will read elsewhere, or even what FTDNA (the testing company I've always recommended for ySTR testing) lays out. All that I can say is that I worked out these principles many years ago and am unaware of anything that has happened since that would require me to change them, and that over the years I've seen signs of others, including FTDNA, coming around slowly to my point of view. Because one of my constant objects in all things is the pursuit of truth and genuine understanding, I will be grateful to anyone who can provide me with new information about these matters, or suggest better ways of thinking about them.

Finally, the fact that there is much in what follows that constitutes implied criticism of FTDNA and its works should not be construed as undermining my admiration and gratitude for what the founder of the company, Bennett Greenspan, and his many helpers have created for us. No one expects the pioneers in a new field to anticipate the best ways of making use of their discoveries and inventions. Rather it is their lot to stake their all on their vision and to hack through and blaze the trail to enlightenment and a brave new world, while it is for those who come after, the users (and particularly the many outstanding ISOGG list activists), to figure out how best to make use of those inventions.

Introduction: Key Concepts and Terms

The most essential term pertaining to ySTR testing and its uses for genealogical purposes is [\(genealogical\) patrilineage](#). The term is my coinage, at least the way I have defined it, but the concept it designates is just as essential to others who use different terms or none. It is mostly one's patrilineage, not one's patrilineal surname per se, that genealogists care about, which is why I've broken down my FTDNA Surname Projects into sets of standalone patrilineage projects—although I also provide an umbrella page for the surname, and the surname project in general.

The [DENNISON Surname DNA Project](#) may serve as an example of the template I've worked out to use for all my projects. Because the individual DENNISON patrilineage projects are each standalone projects, I can use the templates I've created for those pages to create and manage patrilineage projects that are merely satellite to other established FTDNA Surname projects, like the two ALLEN Patrilineage projects I've set up, and I'll be using some of the data in these independent patrilineage projects as illustrations of my principles respecting the interpretation of ySTR test results.

Here are some more essential terms.

The commonly used term MRCA (Most Recent Common Ancestor), or (better in the patrilineage context) [MRCPA \(Most Recent Common Patrilineal Ancestor\)](#), is the most recent patrilineal progenitor of all the members of a set (or subset) of tested male descendants in a patrilineage project. The results of an FTDNA test of a male's ySTR sites is a ySTR [haplotype](#)—a panel of 37, 67, or 111 ySTR marker values. It is because these marker values rarely mutate that the results of an FTDNA 37-marker test can be used to sort descendant males into their particular patrilineage, and it is because a few ySTR mutations can nonetheless be expected to occur over [genealogical time](#) that a patrilineage can sometimes be further divided into sub-branches just on the basis of these test results. Where this is possible, I have coined the term [CCC \(Closer Cousin Cluster\)](#) to represent a subset of descendant haplotypes within a patrilineage project that are more closely related to each other than to any of the other members of the patrilineage, because they all descend from a common MRCPA who lived closer to them in time than the MRCPA they share with all the other members.

The [RPH \(Root Prototype Haplotype\)](#) is my term for the MRCPA's projected ySTR haplotype; the term more commonly used term is "modal" haplotype, or marker value, but this is a misnomer because there's no guarantee that the most common inherited marker or haplotype values in a set of haplotypes are the original ones of the MRCPA, or even provide the best approximation to his haplotype.

Determining an RPH for a set of haplotypes is essential to differentiating mutated marker values from normal (or inherited) ones. There are various ways of doing this, but the most common way, which works well when there are enough haplotypes extended to a particular level (37, 67, or 111), and when it is known from the corresponding genealogies that the ancestries of all of those tested diverged many generations back, is to synthesize the RPH from the most prevalent (or, if you like, the modal) values. However, in cases where there is no overwhelming preponderance of a particular value, the most prevalent value may not be the best choice, and what does one do when there are two equally prevalent values, or three pluralities of values? In such cases, discretion and consideration of both the overall mutational pattern and the associated genealogies is called for.

I will also have much to say below about [TMRCA \(Time back to the MRCA\)](#), whether expressed in generations or years, and [GD \(Genetic Distance\)](#), which in this context of ySTR testing is popularly taken to mean the number of different marker values in the test results of two haplotypes, for example, 35/37, or 105/111. These overt quantitative GD differences are typically equated with the number of mutual mutations that have occurred since the genetic divergence in the ancestral lines of two tested descendants, but as we shall see, counting mutations isn't quite as simple as that. If we take GD at its face value though, TMRCA is simply a function of GD. A probability calculation is applied to a GD difference based on the assumed rates of mutation of the markers involved, or of the marker panel as a whole, and from this the number of generations necessary to produce that many mutations is derived. The number of generations can then be turned into years back to the MRCPA of the two descendant haplotypes by multiplying by the average number of years per generation. Various numbers for this have been used (25 years/generation is conventional in general parlance), but a great deal of empirical evidence points instead to about 34 years, as my paper "[How Long is a Human Generation](#)" argues.

Principle 1: The FTDNA 37-marker test is the one indispensable test for those who want to know for sure to which surname patrilineage they belong; this test is also sufficient to all but rule out the possibility that an NPE (Non-Paternity Event) has occurred in their patriline.

The FTDNA 37-Marker test is both necessary and sufficient for determining a male's patrilineage, and for the majority of testees this is the sole benefit of ySTR testing. How much of a benefit it is depends on the genealogical interactions prompted by the test results, whether these stem from contacts with FTDNA-reported matches, or from the surname or patrilineage projects organized by volunteers around the results. It should be noted though, that even when a person comes up as unmatched, implying that his is a unique one-person patrilineage, there can be great genealogical benefit to knowing that he does *not* belong to certain prominent patrilineages of his surname.

A secondary benefit of determining one's patrilineage is to ascertain whether an [NPE \(Non Paternity Event\)](#) has occurred in one's patriline. And as often as not, only an FTDNA 37-marker ySTR test is capable of doing this, because NPEs so frequently leave no paper trail. NPEs can occur in any generation past, and they typically do occur at rates that vary from about 2-7% or more (until modern times, at least, when rates have soared in many areas). What's more, the chances that an NPE has occurred somewhere up one's ancestral chain are cumulative and amount to between 20-50% for those who can trace their ancestries back 12 or more generations. It's also the case that if one is able to push one's ancestral line back all the way to the period of surname adoption (the 14th century in England (minus Wales), though centuries later in the rest of Great Britain) NPEs are likely to be the rule, not the exception, because the first man to adopt, say, the surname Smith, because he happened to be a blacksmith, is likely to have siblings or cousins who adopted other surnames.

Principle 2: TMRCA and GD comparisons between two haplotypes do not provide genealogically useful estimates of closeness of relationship; they are therefore of little or no value in identifying Closer Cousin Clusters.

This proposition runs counter to what most people believe, and to what FTDNA encourages people to believe. The company has gone to the length of creating a TIP calculator for calculating the TMRCA of pairs of haplotypes (yours and that of one of your matches), expressed as a range of probabilities that your common MRCPA comes in at 1, 2, 3, ... N generations back. The company also makes a point of sorting your matches into closeness of match categories based on GD correspondences of (for the 37-marker test) 32/37 through 37/37. Yet neither of these approaches yield results accurate or reliable enough to be genealogically useful in identifying sub-branches within a patrilineage. And at the same time, the company fails to provide guidance on the analytical techniques that *can* be used to determine which tested members of a patrilineage are more closely related to the other members of what I call their CCC (Closer Cousin Cluster) than they are to all the other members of the patrilineage. I will outline those techniques under Principles 3 and 4 below.

There are many reasons why TMRCA and GD measures don't yield genealogically useful results.

In the first place, the number of marker value divergences between two haplotypes isn't necessarily the same as the number of mutations that have occurred since their ancestral lines diverged.

There are a number of different kinds of mutations, some of which ([multistep mutations](#)) can mimic multiple single mutations, and others (back mutations) that reverse previous mutations back to their original state, thus wiping out all traces that a mutation has occurred.

Then there are the mutations to the [multicopy markers](#), especially the highly mutable DYS464, in which several marker values appear to be changed by a single mutation. There's also a type of mutation called a reClOH event, that can change several multicopy marker values at once by deleting

certain of their values, then effecting a repair of sorts by filling in the missing values from an alternate copy that runs backwards on the same palindromic loop; never mind the technicalities here: what's important is that these reC/OH events, which are quite common (though we have only a vague notion of their frequency) constitute a kind of mutation that can change several marker values at once. Deletions can also occur at other places in the copying of the Y-chromosome, leaving holes (missing marker values).

Determining when these special kinds of mutations have occurred can be quite problematic, and deciding how to account for them—whether to add 0 or 1 or 2 or whatever to the total number of mutations for the haplotype—can fuzz those nice discrete GD comparison numbers and further bloat the already wide range of possibilities when it comes to estimating TMRCA.

In the second place, markers mutate at wildly different rates (some more than 50 time more frequently than others), and besides that, current estimates of these individual marker mutation rates are rather crude, and may be off by as much as 15% or more either way.^[1]

However, these problems in counting mutations, and in applying appropriate mutation rates to the calculation of TMRCA, pale in comparison to this fatal flaw in making pairwise haplotype comparison: **mutations are so rare, and their occurrences so sporadic, that pairwise comparisons even of 111-marker haplotypes over many generations of genetic divergence between two ancestral lineages constitute too small a sample of the mutational process to provide meaningful estimates of closeness of relationship between the descendant haplotypes.**

For example, even a pair of 111-marker haplotypes that diverged from a MRCPA who lived as many as 10 generations ago, provides only $111 \times 9 \times 2 = 1998$ opportunities for mutation, and if one were to make, say, ten such pairwise comparisons, one could expect to find a range of 3-8 mutations separating any given pair, with lesser probabilities of even greater variation. Translating this 3-8 range (over 111 markers and 10 generations) back into TMRCA estimates expressed in years would yield a range of birth dates for one of these pairwise MRCPAs between 1525 and 1800, which makes the estimate genealogically quite useless. And taking such TMRCA estimates seriously is no less misguided than supposing that a 108/111 match is necessarily “closer” than a 103/111 match; these and all the numbers in between are all within the range of normal and expected variability across a set of paired haplotypes.

Let's look at a couple of real-world examples of the meaninglessness of pairwise haplotype comparisons.

In my ALLEN Patrilineage 1 project, the haplotypes of about 20 members have been extended to 111 markers. On the genealogical evidence, which is very extensive and deep, all of these lineages converge backwards to a handful of closely related patriarchs who were born in or immigrated to Virginia between 1680-1710, and probably ultimately to a single immigrant ancestor born between 1620-1650. Although many members of this group remain unconnected to any of the deep ancestral lines, we do know that none of the members of this project shown in the GD matrix converge with any of the others within the last 7 generations, and most converge at about 9 generations on the

¹ It has been claimed that FTDNA's TIP calculator does take into consideration these varying mutation rates from marker to marker, but I also know that, at least in the past, the average mutation rate used by FTDNA for their 37-marker panel was considerably at odds with the rates worked out more accurately by DNA maven John Chandler. Although I myself use these so-called Chandler rates in my own calculations, and even cruder estimated rates for the markers beyond the first 37, it's likely that many of these individual marker mutation rates are wide of the mark, as the underlying mutational frequency data is still accumulating. I do have a fair amount of confidence, however, in the overall average rates of whole marker panels (37, 67, or 111), because there are enough markers in them that probably most of the mis-estimates of individual marker rates cancel out.

average. This large number of highly diverse 111-marker haplotypes has made it possible to calculate a TMRCA for the MRCPA of the whole extended group that is probably accurate to within a generation, or at most two, either way. In fact the predicted birth date of the father of them all based purely on the DNA, which is in the range 1610-1630, dovetails almost perfectly with what the genealogical evidence suggests.

But now look at the range of pairwise GDs in the boxes of [the 111-marker GD matrix for ALLEN Patrilineage 1](#), where the actual differences in closeness of relationship vary only from 8-11 generations. These GDs range from 0 (identical haplotypes) through 12. Even if we exclude the half dozen outliers in this group, the range of GDs still runs from 1-8, and there is little reason to suppose that a pair of these patrilineage cousins (whose lines on average are divergent by about 9 generations, and who are, say, 110/111 are any more closely related to each other than are a pair who are 103/111. At most the odds only slightly favor the former as the closer GD match.

For a second example, one of the members of my ALLEN Patrilineage 2 project has a well-authenticated 12 generation ancestry back to her early New England immigrant ancestor, yet the mutational pattern across the 14 members of this project who have tested to 111 markers clearly shows that her haplotype has experienced no mutations in all that time. The odds of this happening, given the known number of generations and the mutation rate for the 111-marker panel, are just 2%, but long odds cases do, by definition, turn up from time to time.

For a contrary example, consider one of the members of my RATHBONE DNA surname project who has now tested himself, his father, and his brother through 67 markers. As expected, the haplotypes of father and brother are identical, yet the subject himself has experienced two mutations in just one generation--again, there's only about a 2% chance of that happening.

Finally, FTDNA's TIP calculator page itself recognizes one other major problem with the TMRCA estimates the calculator turns out. Suppose you use TIP to compare your haplotype with one of your closest matches, who is 111/112 with you. The TIP calculator says, not very helpfully, that there's a cumulative 55% chance that your common ancestor goes back no more than 2 generations, an 85% chance of no more than 4 generations and a 96% chance of no more than 6. As you are trying to guess where in this wide range of generations your common ancestor is located, thinking, perhaps that the 55% (or 2 generations) would be the best guess, suppose that you happen to learn that in fact the relationship goes back exactly 6 generations, not 2 (I'm basing this example, incidentally, on a real case in one of my projects--in fact, the first case that came to mind).

The TIP calculator looks a bit silly here, estimating that the two subjects in this actual case were 1st cousins, when in fact they are 5th cousins, but as I say, FTDNA has made provision for cases in which the genealogy of the two subjects is known well enough that one can be quite sure that they have no common ancestor back a certain number of generations. In this case, where the exact number of generations back to the MRCPA is known (here 5), entering that parameter on the TIP page, produces a new range of TIP probability estimates: a 47% probability of a convergence at the 6th generation, a 72% chance at the 7th, 85% at the 8th, and so forth. But ironically, even though TIP has here been fed the actual answer to the question asked, it still provides only a range of probability estimates that are still wide of the mark. TIP simply isn't able to provide a genealogically useful estimate for then the MRCPA of a pair of haplotypes comes in.

Although individual pairwise GD comparisons are of little or no genealogical value, constructing a matrix of *all* the pairwise comparisons across a large patrilineage can shed light on the overall depth and shape of the patrilineage yDNA tree as a whole, and sometimes it can at least suggest where the Closer Cousin Clusters (CCCs) are to be found. When one arranges the members of a large GD matrix according to their (on average) GD closeness to all the other members collectively, this

sometimes results in several different clusters of low numbers (shaded green), each suggestive of an independent CCC. The [111-marker GD matrix of the ALLEN Patrilineage 2 project](#) provides an example of this: the last three haplotypes listed appear, just from the chart, to constitute a small CCC independent of the remaining members of the project, and in fact, this is borne out both by their genealogies (which indicate that they descended from a different 17th century immigrant ancestor), and also by the more effective methodology for determining CCCs that I will explain next.

Principle 3: Analysis of Mutational Patterns across a patrilineage can identify Closer Cousin Clusters (CCC), though only under special conditions of knowledge that do not obtain for most current sets of patrilineage members.

The ALLEN Patrilineage 1 project cited above is the kind of project for which CCCs can be identified, as it has the chief desiderata: many haplotypes of diverse lineages that have been extended to 111 markers, backed up by solid and deep genealogical knowledge. In fact this project has 19 such haplotypes extended to 111 and is the largest that I've worked with or studied to date. How many is the "many" 111-marker haplotypes that are needed to identify CCCs? I'd say at least 5 or 6, and for 37 markers, 7-10—and this is presuming that all of the haplotypes in question are known to diverge from each other over at least 6 generations. With fewer than these numbers it becomes problematic to determine the composition of the RPH, and therefore to identify which values are mutations, and which the normals, and it is upon the mutational pattern that inferences to CCCs are predicated.

The additional requirement, the need to be sure that the descendant haplotypes have no common patrilineal ancestor within at least the last six or so generations implies a fourth principle of ySTR DNA interpretation:

Principle 4: The identification of and exploitation of Closer Cousin Clusters depends more on genealogical research than on ySTR DNA testing.

Beyond the sorting of haplotypes into patrilineage (which is conclusive by itself according to Principle 1), ySTR DNA testing can at most provide only *guidance* toward the most fruitful avenues of genealogical research: it cannot substitute for that research. When it comes to determining CCCs, DNA-based theories must generally yield to the genealogical evidence when there are conflicts, though in extreme cases, where the DNA probabilities point the other way with a 95-98% confidence level, the genealogical evidence and the conclusions drawn therefrom need to be scrutinized very critically.

Most of the time, however, the mutational patterns in the DNA that appear to demarcate CCCs confirm, or are at least consistent with what the genealogical evidence suggests, and the two modalities of evidence mutually strengthen each other.

To return to the methodology for analyzing the mutational patterns that characterize CCCs, in light of Principle 4 the set of (extended) haplotypes used as the basis for the analysis needs to be pruned of all known closer cousins (or patrilineal relatives) from recent generations, both to achieve the six or so generations of known separation between cousins, and also (as a preliminary) in deriving the RPH that serves as the standard for determining which marker values are the mutated ones, and which the normal values. In general, all but one of these recent clusters of known closer cousins need to be weeded out, leaving that one to represent the rest of his recent lineage. A corollary of this rule is that when it comes to ySTR testing, there is little or no benefit to testing known close patrilineal relatives—ones' father, brothers, or close cousins—unless one simply wants to rule out the possibility that one of them may have experienced an NPE in his line, and for that purpose, the lowest resolution ySTR test that FTDNA offers should suffice, and that would seem to be the 37-marker test (too bad, since FTDNA used to offer a much cheaper 12- or 25-marker test).

Principle 5: Ultimately, the value of ySTR DNA testing for genealogical purposes depends on the depth and quality of evidence that your patrilineage cousins are able and willing to share.

The first requirement here is that the tests that have been run to produce at least a 37-marker ySTR haplotype which turns up as a match to a particular patrilineage, have an active and knowledgeable genealogist behind it, whether that be the patrilineal descendant male who actually tested, or his sponsor or genealogical representative.

The second requirement is that this genealogist remains prepared to respond to inquiries at the email address he has supplied to FTDNA, or to the surname or patrilineage projects s/he may have joined that have been organized around FTDNA ySTR test results.

Unfortunately, all too many FTDNA customers order tests without understanding their limitations, or what they need to bring to the table to take advantage of them; many are looking for a genealogical shortcut or a magic talisman that will somehow get them past their brick wall. However, unless they are fortunate enough to be able to hook up with a robust and genealogically activist patrilineage project, the most that they can hope for is that one of their matches will (a) respond to their email query; and (b) be more genealogically knowledgeable about their particular line than they are. Alas, all too often, the few testees who are very genealogically knowledgeable about their surname patrilineage have over time derived so little benefit themselves from their tests (precisely because they know more than their matches), that they don't bother to respond, or to keep their email address up-to-date.

Another reason FTDNA-reported matches often fail to respond is that in the common case where the interested genealogist has tested, not themselves, but a surrogate male who bears the surname that they're interested in, the one email address that FTDNA presents to their matches is that of the surrogate, and not the genealogist—and many if not most surrogates, not having much interest in their own genealogy, fail either to respond or to pass on the contact to their genealogical representative.

This comes about, because when FTDNA initially solicits these email addresses from the customer, it merely asks him/her to list one or more email addresses in any order, first in a field associated with the customer herself and with her physical address, and then in a second blank associated with the surrogate and his physical address, without explaining which of these addresses will be used as the “primary” email address for purposes of genealogical contact, or even making it quite clear which of the two physical addresses is to be used in shipping the test kit. I suppose that the primary email address would be the first listed in the customer blank, but since this first blank is associated with a physical mailing address, which the customer may be forgiven for assuming that this first physical address is the one for the test kit, s/he may also assume that the first email address listed should correspond to that physical address, and also be that of the surrogate. All of this ambiguity could be avoided by clarifying the entry form, and dedicating a field to “Email address to be contacted at by your test result matches?” And while they are at it, it would be also be desirable to expand this contact information to include more than one email address, lest one become obsolete over time.

The most important factor regarding matches, though, is simply the quality and depth of their genealogical research. One advantage of independent projects organized around the patrilineage is that instead of just providing a few pairwise matches, the contact information for all of the principal genealogists interested in the line can be provided in a project directory that is also available to other researchers interested in the patrilineage who are as yet untested, but have found the web page through browsing. And in general, the best way to take advantage of ySTR DNA testing for genealogical purposes is to organize in or participate in such patrilineage-oriented projects.

The CCC Analytical Method, applied to a particular patrilineage

Let's look at another example of a patrilineage for which CCC Analysis has proved fruitful—the [DENNISON Patrilineage 1 Project](#). In this large patrilineage of 17 members, the haplotypes of 11 have been extended to 111 markers (here's [the project 111-marker GD matrix](#)). These lineages have been very thoroughly researched, and most members can trace back to their immigrant ancestor. Thus, we can also say of this set of 11 extended haplotypes that we know that their lines don't converge for the last six generations, and that the average distance back to the MRCPA is probably about 9 generations.

The general idea in CCC analysis, however, is to study, not the GD matrix, but the mutational pattern shown by [the project haplotype chart \(or table\)](#). In this table of marker value results for each haplotype, mutated marker values are those identified by the colored squares, and the different colors represent different types of mutations—as defined by the color-coding glossary in the text at the bottom of the table. It will be noted that I've grouped the haplotypes together into clusters in the light both of the mutational patterns suggested by the DNA data itself, and also from my knowledge and suppositions regarding the actual ancestral relationships among these subjects.

Definitive identification of CCCs depends on finding distinctive mutations that all the members of a particular cluster share, which they inherited from the MRCPA of their cluster.

But this formulation begs several questions.

First, there is the problem discussed briefly above: how do we determine which marker values are the mutated ones, and which the normal — the values that comprise the RPH we wish to construct as our best guess for the composition of the haplotype of the MRCPA ancestor of all members of the patrilineage?

As noted above, in most cases, one can simply use the preponderant value across all the haplotypes, but where the preponderance is slight (with one value having only a simple majority over the other, or where the subsets of values are of equal size), and/or where there are more than two marker values, other principles of analysis must be brought into play. Where there are, say, three values, the most likely original MRCPA value is the middle one, since the mutational process is generally suppose to run either way, in roughly equal proportions, either adding or subtracting a number to the original marker value. However, certain directional biases seem to be emerging from the so far minimal mutational data favoring either gaining or losing a “[repeat](#)” (in the technical terminology), so we may need to keep an eye on this possibility going forward.

Where the balance is fine between two marker values, I take into account what I know of the genealogy, and of the likely genealogical clusterings and weightings within the particular set of haplotypes involved. Thus, if it appears that 6 of 111 haplotypes have marker value A, and 5 value B, but I know or believe that most of those who have value A look like the makings of a Closer Cousin Cluster either because their haplotypes are similar otherwise, or because their ancestries seem to converge short of the MRCPA of all, while the smaller set of 5 haplotypes shows greater diversity either of DNA or genealogy or both, I may chose the minority value B as the normal value, and use it in constructing the RPH. As noted above, drawing on the genealogical as well as the DNA evidence introduces an element of circularity if our aim was to reason out relationship strictly from the DNA evidence, but in view of Principle 4, our DNA-analytical goal here must be the more modest one of just looking to the DNA evidence for suggestive, or confirmatory patterns to support whatever we can learn from genealogical research.

It's not always going to be possible to pick the correct marker value as the mutated one, and it may be necessary to change one's initial picks as more, and particularly more diverse, haplotype accrue to

the project. The reason for this inherent uncertainty about which values are the mutated ones, is analogous to uncertainty bedeviling pairwise TMRCA: in both cases, the sample size is too small.

The first problem one encounters in CCC analysis, correctly ascertaining the RPH, and thus identifying the mutated values, can be expected to evanesce as projects grow in size. However, there's a second type of problem may actually increase as more haplotypes come into play.

This biggest problem with using shared mutations to identify CCCs is that sometimes two haplotypes may share a mutation, not because they inherited it from a common ancestor, but because the mutation occurred independently in two or more sub-branches of the patrilineage that aren't closely related to each other.

Unfortunately, there is no definitive way to determine whether a shared mutation was inherited rather than generated independently a second or third time within the same set of haplotype. However, we can look to the widely varying mutabilities of the individual markers to assess the corresponding probability that a particular marker may have mutated more than once in the same patrilineage.

The Mutational Quirks of Certain Markers

As it happens, the very most mutable markers in the FTDNA panels (DYS449, 464, the CDYs, and 710 and 712 in the 111-marker panel) are quite likely to have mutated independently more than once across a fairly large set of haplotypes (say 7-10), and there are many other fast-mutating markers that are suspect.

I roughly indicate the relative mutability of individual markers in the columnar marker headings of my haplotype charts— color coding the faster mutators red (the fastest are the deepest reds); the average mutators with the background text color (a dark reddish brown); while the notably slow mutators are color-coded blue. Naturally, I use the more exact (but still only approximately known) numeric probabilities for each marker in my probability calculations.

Thanks to the ambiguous possibilities here, **where shared mutations may or may not be inherited, reasonably reliable identification of a CCC from the DNA evidence alone typically depends on finding at least one a shared average or slow mutator, or more than one fast mutating marker.** Of course, where there is also strong genealogical evidence identifying a CCC, lesser DNA evidence may be used to reinforce that identification.

Finally, in mutational assessment, a special word is called for with respect to two sets of markers. The pair of CDY markers in the 1-37 panel mutate so frequently that they are best ignored in these analyses, except, perhaps as confirmatory grace notes. The probability that one of these will mutate in a single father-to-son generational transmission is about .03531, thus the probability of it mutating more than once in a single haplotype over, say, 10 generations is 7%. And if one is analyzing a set of 10 haplotypes, one can positively expect more than one such mutation to each CDY, and once in a while a second mutation in the same haplotype that's actual a back mutation to it's original value.^[2]

When the **DYS464** multicopy marker, which typically comprises four component values (and sometimes several more) mutates it often changes more than one of its component values at once, and "calling" the values of this marker even has an element of subjective interpretation on the test bench. Without getting into the technicalities of why this is so, **DYS464** is best treated as a single highly

² From the point of view of analysis, the instability of these CDYs isn't as bad as it seems: because a marker can mutate up or down the chances that the same CDY is a different haplotype will mutate in the same direction as a previous one is just half this ultrafast mutation rate; on the other hand, though, a mutated CDY marker has an equal, appreciable, chance of mutating back to its original value, leaving no trace that it ever mutated at all.

mutable marker, with a per generation mutation probability of .02264—not far behind the CCCs (the next most mutable marker is DYS710, at .01570). Offsetting this, the way DYS464 is normally evaluated, back mutations rarely occur.

Precisely because DYS464 comprises 4+ marker values and is so mutable (two thirds as mutable as the CDYs) we can't afford to ignore it in our haplotype cluster analysis. And FTDNA offers an optional test called DYS464X that eliminates the ambiguities in the evaluation of DYS464 and more often than not reduces it to 2 marker values that are just as solid as any of the single-copy markers. I've therefore generally recommended, whenever a patrilineage has turned up several different DYS464 values according to the default test methodology, that at least certain of the project members order the additional DYS464X test, so that we can figure out what is really going on with this important marker. Unfortunately, FTDNA seems to have decided to bundle this often valuable test with re-evaluations of other multi-copy markers that aren't of such value, with a higher price on the bundle, so I expect to use more circumspection in recommending this test to my project members in the future.

With the above principles and caveats in mind, you may wish to read through [my detailed CCC analysis of DENNISON Patrilineage 1 haplotypes](#) as an extended example of what can be done with a large set of extended haplotypes. Since the analysis is specific to the mutational pattern across these 11 extended haplotypes, it would be well while reading this material to bring up [the corresponding project haplotype table](#) in a separate window for reference. The table scrolls to the right, allowing one to see which haplotypes have extended to 111, and showing the mutations that have occurred in that upper band.